

COLONOSCOPY INSTRUCTIONS ARH (page 1/2)

Dr. Yuanyuan Chen
110-3113 272 St, Aldergrove, BC

604-381-1800
email: yychenclinic@gmail.com

Date of procedure: _____ **Check-In time:** _____

Procedure location: **Abbotsford Regional Hospital General Daycare Unit**

32900 Marshall Road, Abbotsford

YOU MUST PHONE OUR OFFICE (604-381-1800) TO CONFIRM YOUR APPOINTMENT 14 DAYS BEFORE YOUR APPOINTMENT. WE REQUIRE 2 WEEKS NOTICE FOR CANCELLATION OR RESCHEDULING; OTHERWISE THERE WILL BE A CHARGE OF \$500.

Stop all blood thinners prior to colonoscopy unless otherwise directed by your family physician or specialist. Consult your family physician or specialist to see if you will require bridging therapy while off blood thinners.

- stop plavix/clopidogrel, warfarin/coumadin, ticagrelor/brilinta 5 days prior to colonoscopy
- stop apixaban/eliquis____days prior to colonoscopy
- stop dabigatran/pradaxa____days prior to colonoscopy
- stop rivaroxaban/xarelto____days prior to colonoscopy

If you take medications for diabetes, please consult your GP for directions for the day of the colonoscopy.

You require a driver to bring you home from the hospital. Due to sedation, you are considered legally impaired for up to 24 hours post procedure. You CANNOT drive, take taxi or bus home post colonoscopy.

If you need an interpreter, you must provide one for yourself and they must attend both check in and discharge with you.

FYI: inpatients needing urgent procedures take priority over outpatients. Sometimes outpatients need to be rescheduled to accommodate urgent inpatient procedures.

You will be in the hospital for approximately 2.5-3.5 hours from your check in time.

COLONOSCOPY INSTRUCTIONS ARH (page 2/2)

Dr. Yuanyuan Chen
110-3113 272 St, Aldergrove, BC

604-381-1800
email: yychenclinic@gmail.com

The doctor's ability to visualize your colon depends on the thorough bowel preparation. Please follow instructions closely. You will need to repeat this procedure if your bowel preparation is inadequate.

7 days before colonoscopy:

Stop all anti-inflammatories (e.g. advil/motrin/ibuprofen, aleve/naproxen, celebrex/celecoxib, cambia/diclofenac, Toradol/ketorolac), and oral iron supplements. Please check multivitamins for iron as well.

- Purchase 1 bottle of Colyte 4L or Peglyte 4L (no substitute).

5 days before colonoscopy:

- Check the first page for instructions on stopping blood thinners.
- Stop eating foods that contain seeds (e.g. blueberries, strawberries flax seeds, etc), popcorn, corn, granola, shells, multigrain bread, nuts, stringy/fibrous fruit & vegetables.
- Eat a low fiber diet (e.g. plain white bread (no crust), white rice, applesauce, potatoes/yam/sweet potatoes (no skins), steamed/cooked carrots, cream of wheat, eggs, lean meats and well-cooked porridge).

1 day before colonoscopy:

- Prepare the 4 liters of Colyte or Peglyte as directed on the jug. Keep in fridge as it is best taken cold rather than at room temperature.
- Start clear liquid diet. Have a variety of clear liquids to replace electrolytes. Clear liquid examples are clear broth, water, clear pulp free juices, apple juice, white grape juice, white cranberry juice, tea & coffee with NO milk or cream, clear GingerAle or Sprite, clear Gatorade, jello (only yellow, green). No red, purple drinks or jello of any kind. No alcohol. No milk, soy or almond milk products.
- Stay close to a toilet once you start taking the bowel preparation as it will cause diarrhea.
- At 4pm, drink 2 liters of the Colyte or Peglyte within 90 mins (drink 1 glass, 250mls, every 10 minutes). It is often better tolerated if well chilled in advance and taken through a straw.
- 5-6 hours before your arrival time, drink the remaining 2 liters of the Colyte or Peglyte whining 90 minutes.

The day of colonoscopy:

- Stop having clear fluids at least 3 hours before your arrival time.
- You may continue to take your usual medications with a sip of water unless otherwise directed by your family doctor or specialist. If you take medications for diabetes, please consult your GP for directions.
- Bring a list of your medications, care card with picture ID to the hospital.
- Do not wear perfume to the hospital.